

A Review on Diabetes

Amit Kumar Agrahari¹, Abhinav Shukla², Diwakar Shukla³, Abhishek Dubey⁴, Aditya Kumar⁵, Kunal Chaurasiya⁶,
Rajneesh Gupta⁷, Saloni Vishwakarma⁸

Praduman Singh Shikshan Prashikshan Sansthan Pharmacy College, Basti, India

Abstract--- Diabetes mellitus is a general term for heterogeneous disturbances of metabolism for which the main finding is chronic hyperglycaemia. The cause is either impaired insulin secretion or impaired insulin action or both. High blood glucose levels are symptomatic of diabetes mellitus as a consequence of inadequate pancreatic insulin secretion or poor insulin-directed mobilization of glucose by target cells. Diabetes mellitus is aggravated by and associated with metabolic complications that can subsequently lead to premature death. This review explores diabetes mellitus in terms of its historical perspective, biochemical basis, economic burden, management interventions along with the future perspectives.

Key Points-- Diabetes mellitus, Types

I. INTRODUCTION

Diabetes mellitus (DM) is the most common endocrine metabolic disorder which caused by improper regulation of insulin secretion or insulin action. Deficiency of insulin leads to chronic hyperglycaemia with disturbances of fat, protein and carbohydrate metabolism. [1-4]

Long term diabetes cause tissue or vascular damage and leads some dangerous complications like neuropathy [5,6], nephropathy[7,8], retinopathy [9,10], cardiovascular complications (11,12) and ulceration. [13,14] First diabetic classification and diagnostic criteria was given by World Health Organization (WHO) in 1965[15] and the National Diabetes Data Group (NDDG) in 1979. [16]

Symptoms:- Blurred vision, polyurea, polyphagia, polydipsia, weight loss, constipation, cramps, thirst, hunger, weakness, [17,18] and candidiasis are common symptoms for both type-1 and type-2 DM. [1] Apart from that the patients with Type-1 DM have both microvascular problems [5-10] and macrovascular diseases such as heart, peripheral vascular and coronary artery diseases [11,12] while patients with Type 2 DM have high risk of large vessel atherosclerosis commonly associated with obesity, hypertension, hyperlipidaemia. [11,12,19,20] Most patients are die with type-2 diabetes due to cardiovascular complications and end stage renal disorder [9-12]




Fig-1 Diabetes

II. TYPES

There are several types of diabetes mellitus but two main types are type-1 and type-2. [21,22] On the basis of aetiology type-1 is describe as IDDM (juvenile-onset diabetes) and type-2 describes as NIDDM (maturity-onset).




Fig- 02-Types of diabetes

Type-1 diabetes:- It occurs in those patients who have no or little endogenous insulin secretory capacity and require insulin therapy for survival. Type-1 diabetes is further divided into two categories; Type-1a:- which is characterized by presence of islet cell antibody (ICA), anti-glutamic acid decarboxylate (anti-GAD) and insulin antibodies (IA-2) that identify the autoimmune process with β -cell destruction. [22,23] and is caused by immunological destruction of pancreatic β cells and leads to insulin deficiency and other is type-1b; which is idiopathic and have no autoimmunity. [23,24]

Type-2 diabetes:- Globally 5-7% of the world's population [25, 26, 27] and in Western countries up to 7% of the population is affected by type-2 diabetes. [27, 28] Urban areas have high incidence of type-2 diabetes than in rural areas. [26, 29, 30] Type-2 diabetes is associated with obesity, decreased physical activity, heredity and characterized by the disorders of insulin secretion and insulin resistance. [31] It is common in individuals over the age of 40 [19, 20] and is usually controlled through dietary therapy, exercise and hypoglycaemic agents. [32,33]

Gestational Diabetes (GD) mellitus:- Gestational diabetes is occurs during second or third trimester of pregnancy. [34] It occurs in about 4% of all pregnancies. GD Patients have 30% to 50% chance of developing DM, usually type 2 DM.

Other types of diabetes mellitus include genetic defects of the pancreatic β cell or in insulin action pathways (insulin receptor mutations or post-receptor defects) [35], Endocrinopathies producing insulin counter regulatory hormones (e.g., Cushing's syndrome, acromegaly), disease of the exocrine pancreas (e.g., Pancreatitis, pancreatic reaction, or cystic fibrosis) may produce DM. (36) several medicines such as niacin, glucocorticoids, pentamidine, and a-interferon may also lead to DM.[37]

Epidemiology:- In 2011, 4.6 million deaths occurs by DM and estimated that 366 million people are suffering from diabetes mellitus in 2011 which may risen to 552 million in 2030. [38] In 2030, 439 million people would have type 2 DM and the causes of type 2 DM varies by environmental and lifestyle risk factors. [39] The incidence of DM is common in adults of which type 2 DM is becoming prominent and will increase in the next two decades in developing countries where most of the patient are aged between 45 and 64years. [40]

III. DIABETES IN INDIA

In 2010 approx 285 million people worldwide (6.6%) have diabetes and most of the diabetic patients are belonging to the age between 20-79years which will increase up to 438 million people (7.8%) in 2030. According to Diabetes Atlas 2006 which was published by International Diabetes Federation, around 40.9 million people have diabetes in India which is expected to be rise up to 69.9 million. [41] In 2000, India (31.7 million) topped the world with the highest number of people with diabetes mellitus followed by China (20.8 million) with the United States (17.7 million) in second and third place respectively. [10,14] Diabetes is potentially epidemic in India with more than 62 million diabetic persons currently diagnosed with this disorder. [42, 43]

IV. FREE RADICALS

In human body, during manufacture of plastics, ageing of paints and the combustion of fuels produce Free radicals in a wide range of chemical and biological systems. In living organisms, free radicals and other 'reactive species are controlled by a group of antioxidant defences, which reduces oxidative damage to bio molecules. In human disease, the balance of 'oxidant-antioxidant' is tilted in the favour of reactive species, so that oxidative damage levels increase.

In some diseases, this may contribute to tissue injury, provide growth for therapeutic intervention with rationally designed antioxidant drugs. [44, 45] Free radical can be defined as an atom or molecule containing one or more unpaired electrons, an unpaired electron being one that is alone in an orbital and is capable of independent existence. [46] Free radicals are involved in many pathological conditions such as many types of diabetes, neurodegenerative diseases, cardiovascular diseases (CVDs), cancer, cataracts, asthma, rheumatoid arthritis, inflammation, burns, intestinal tract diseases, progerias and ischemic and post-ischemic pathologies. [47]

V. SOURCES OF FREE RADICALS

The ROS can be produced from either endogenous sources such as mitochondria, peroxisomes and endoplasmic reticulum, where the oxygen consumption is high or exogenous sources.

Mitochondria:- Mitochondria is capable of producing intracellular ROS. The superoxide radicals are produced from two major sites in the electron transport chain, namely NADH dehydrogenase (complex I) and ubiquinone cytochrome “c” reductase (complex III). The generation of superoxide is non-enzymatic and therefore higher the metabolic rate, the greater is the production of the ROS (48). Monoamino oxidase, aketoglutarate dehydrogenase, glycerol phosphate dehydrogenase and p66shc are the other mitochondrial components which contribute in formation of ROS. [49]

Peroxisomes:- In peroxisomes the respiratory pathway involves in transfer of electrons from various metabolites to the oxygen leads to H_2O_2 formation [50], but is not coupled to oxidative phosphorylation to produce ATP, only free energy is released in the form of heat. beta-oxidation of fatty acids is the major metabolic process of producing H_2O_2 in the peroxisomes. H_2O_2 , $O_2^{\bullet-}$, $OH^{\bullet}$ and $NO^{\bullet}$ are other free radicals which are produced in peroxisomes. There are different peroxisomal enzymes such as acyl CoA oxidases, D-amino acid oxidase, L-alpha-hydroxy oxidase, urate oxidase, xanthine oxidase, D-aspartate oxidase are capable to produce different ROS. [51]

Endoplasmic Reticulum:- Diamine oxidase and enzymes of endoplasmic reticulum such as Cytochrome p-450 and b5 contribute to the formation of ROS. The other endogenous sources of ROS include prostaglandin synthesis, auto-oxidation of adrenalin, phagocytic cells, reduced riboflavin, FMNH₂, FADH₂, cytochrome P 450, immune cell activation, inflammation, mental stress, excessive exercise, infection, cancer, aging, ischemia etc. [52]

Oxidative stress:- Oxidative stress is the situation of a severe imbalance between production of reactive species and antioxidant defence. In 1968, Irvin Fridovich discovered superoxide dismutase (SOD) which is a very unique enzyme that catalyzing the transition of $O_2^{\bullet-}$ into H_2O_2 . [53,54] Then few years later, Chance and coauthors reported that mitochondria are the key producer of $O_2^{\bullet-}$ in cells [55] and has been found that free radicals are produce in biological system and their concentrations are regulated by enzymatic mechanism. Later it has been proved that non-radical products such as H_2O_2 or hypochlorous acid, which are also powerful oxidizing agents, participated in free radical reactions same as free radicals (56). Radicals and the non-radicals species are collectively known as “reactive oxygen species” (ROS) where as Nitric oxide ($NO^{\bullet}$) and peroxynitrite were also interact with ROS, and all of these species were known as RONS which is produced by immune system [57] but excessive production of RONS would damaged almost all classes of biomolecules, such as lipids [58] proteins [59] and DNA. [60, 61] The term “oxidative stress” was introduced in 1970s and 1980s to describe these deleterious processes. Helmut Sies defined oxidative stress as; it is an imbalance between oxidants and antioxidants in favor of the oxidants, potentially leading to damage. [62] Induction of oxidative stress is very important for immune system as it play significant defense mechanism against bacteria. [63]

After observation of oxidative damage reactions of biomolecules shows that primary RONS such as $O_2^{\bullet-}$, H_2O_2 or $NO^{\bullet}$, in most cases reversibly react with the target molecules and oxidative damage is primarily associated with secondary RONS, such as $^{\bullet}OH$, $ONOO$ and $HOCl$ (64-66). There are two major reactions leading to the formation of toxic RONS are: (i) the Fenton reaction between ferrous ions and H_2O_2 yielding $^{\bullet}OH$; and (ii) the reaction of $O_2^{\bullet-}$ with $NO^{\bullet}$ yielding $ONOO$. Iron ions can directly react with organic peroxides, inducing lipid peroxidation and presence of iron and copper ions involve in double-strand breaks of DNA. [67] Oxidative damage to proteins is regularly associated with the reaction between amino acids and $ONOO$ which forms nitrated amino acids like nitrotyrosine. [68]

VI. RONS

ROS (reactive oxygen species):- oxygen derived radicals are most important radical species which are generated in living system. Molecular oxygen is a radical and has a special electronic configuration. Addition of one molecule to dioxygen forms superoxide anion radical [69] and it is generate mainly in a cell of mitochondria. [70]

Superoxide anion are generated through metabolic processes or by physical irradiation which called as primary ROS and these ROS further interact with other molecule to generate secondary ROS.[71] Both complex 1 and complex III electron transport chain are involved in producing superoxide and when they are in anionic form, it is more charged and easily crosses the inner mitochondrial membrane. Recently, it has been confirmed that complex I dependent superoxide is completely released into matrix.[72] The superoxide radicals participated in the Haber- weiss reaction which combines a fenton reaction and the reduction of Fe^{3+} by superoxide yielding Fe^{2+} and oxygen [73] Hydroxyl radical is very dangerous because of its high reactivity and half life of approximately 10⁻⁹s in vivo. Hydroxyl radical is the neutral form of hydroxide ion. [74] According to fenton reaction the most flexible hydroxyl radical is produce when Mn^{+} is iron, chromium, cobalt and copper[75] Organism which have higher amount of iron (hemochromatosis, b-thalassemia, hemodialysis) contain large quantity of “free available iron” and are transported into an intermediate, labile iron pool which represents a steady state exchangeable and readily chelatable iron compartment. [76] Other reactive radicals which are produced by oxygen in living system is peroxy radicals and $HOO^{\bullet}$ is the most simple form of

peroxy radical, called hydroperoxy radical [77] Hydroperoxy radical begin the fatty acid peroxidation by two pathways:- fatty acid hydroperoxide independent and LOOH dependent ways. [78]

RNS (reactive nitrogen species):- Nitric oxide is an important reactive radical that act as oxidative biological signalling molecule in different physiological processes such as defence mechanism, blood pressure regulation, immune regulation, neurotransmission and smooth muscle relaxation. [79] $NO^{\bullet}$ is soluble in both liquid and lipid medium and readily diffuse through cytoplasm and plasma membranes Half-life of $NO^{\bullet}$ is only few seconds in an aqueous environment and has greater stability in an environment with low oxygen concentration. [80] NO reacts with oxygen and water to form nitrate and nitrite ions. Excess production of reactive nitrogen species called nitrosative stress [81,82] which may lead to nitrosylation reaction and alter the protein structure and inhibit their normal function. During inflammation process, cells of immune system produce both super oxide anion and nitrogen oxide during oxidative burst and readily react together to produce oxidatively active molecule such as peroxynitrite anion which is a very potent oxidising agent that can cause DNA fragmentation and lipid oxidation.[83]

Table.1. Different types of ROS and RNS produced in the cell

Reactive Oxygen Species (ROS)

Radicals:		Non-Radicals:	
$O_2^{\bullet-}$	Superoxide	H_2O_2	Hydrogen peroxide
$OH^{\bullet}$	Hydroxyl	$HOCl^{\bullet}$	Hypochlorous acid
$RO_2^{\bullet}$	Peroxy	O_3	Ozone
$RO^{\bullet}$	Alkoxy	1O_2	Singlet oxygen
$HO_2^{\bullet}$	Hydroperoxy	$ONOO^{\bullet}$	Peroxynitrite

Reactive Nitrogen Species (RNS)

Radicals:		Non-Radicals:	
$\text{NO}\cdot$	Nitric Oxide	ONOO^-	Peroxynitrite
$\text{NO}_2\cdot$	Nitrogen dioxide	ROONO	Alkyl peroxynitrites
		N_2O_3	Dinitrogen trioxide
		N_2O_4	Dinitrogen tetroxide
		HNO_2	Nitrous acid
		NO_2^+	Nitronium anion
		NO^-	Nitroxyl anion
		NO^+	Nitrosyl cation
		NO_2Cl	Nitryl chloride

Relation between oxidative stress and diabetes:-

Immune response of the body such as response against self-cells and self-biomolecules is directly linked with all 3 types of diabetes so any disturbance in immune response can cause any type of diabetes. In many other studies it is proved that the auto-reactive CD8⁺ T cells are able to cause type-1 diabetes where as immune cells destroyed the beta islets of Langerhans and cause type-1 diabetes. [84,85] Same as Type 2 and Type 3 diabetes are connected with immune response. It was hypothesized that the non-specific response of the immune system may produce diabetes and found that the patient with diabetes have association of interleukin-6 (IL-6) and C-reactive proteins (CRP). [86] A review showed that increase fat content in food leads to an increase in CRP and IL-6 and ultimately increasing the risk of diabetes. Free radical peroxidation of lipids causes destruction of enzymes and receptors activity which are bound to the membrane and local damage to cell membranes, lastly injured the organs. False immune system cause diabetes and therefore lipid peroxidation is takes place. Stress can arouse hypoglycemia, hyperglycemia or have no affect on glycemic condition in diabetes. [87] In obese person, the adipose tissue becomes saturated with fat, adipocytes starts secreting low level of tumor necrosis factor- α , and stimulates preadipocytes for production of monocyte chemoattractant protein-1.

High lipid content of adipose tissue also triggers oxidative damage. [88] In a study it has been clearly shown that the immune reactions involve ROS, causes lipid peroxidation. [89] and observed excessive production of free radicals in both Type-1 and Type-2 diabetes.[90] In hyperglycemic condition oxidative stress is increased which cause peroxidation of cellular membrane lipids and enhance oxidative modification of DNA and amino acids [91] In type-2 diabetic patients oxidative modification of Phospholipids is takes place which leads to increase peroxide lipids. [91]

VII. CONCLUSION

The global burden of diabetes is increasing worldwide as it is a costly disease for developing economies of the world. To reduce the pandemic of type 1 and type 2 diabetes and its effects on lives and economies worldwide, it is necessary to have an improved understanding of its etiology, pathogenesis and pathophysiology to focus therapeutic and research efforts appropriately. A coordinated multidisciplinary approach is needed that involves scientists, public health practitioners, educators, clinicians and diabetics, with support from government authorities and nongovernmental organizations to reduce the incidence of diabetes significantly

REFERENCES

- [1] Kumar PJ, Clark M. Textbook of Clinical Medicine. Pub: Saunders (London), 2002, pp 1099-1121,
- [2] Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. Report of the expert committee on the diagnosis and classification of Diabetes Mellitus. *Diabetes Care* 1997; 20: 1183-1197.
- [3] Beverley B, Eschwège E. The diagnosis and classification of diabetes and impaired glucose tolerance. In: Textbook of Diabetes 1 Ed: John C Pickup and Gareth Williams Third edition; Chapter 2, 2003, pp 2.1-2.11.
- [4] Lindberg G, Lindblad U, Melander A. Sulfonylureas for treating type 2 diabetes mellitus. *Cochrane Database Systemic Reviews* volume 3, 2004.
- [5] Seki M, Tanaka T, Nawa H . Involvement of brain-derived neurotrophic factor in early retinal neuropathy of streptozotocin-induced diabetes in rats: therapeutic potential of brain-derived neurotrophic factors for dopaminergic amacrine cells. *Diabetes* 2004; 53: 2412-2419.
- [6] Moran A, Palmas W, Field L . Cardiovascular autonomic neuropathy is associated with microalbuminuria in older patients with type 2 diabetes. *Diabetes care* 2004; 27: 972-977.
- [7] Bearse MA Jr, Han T, Schneck ME. Local multifocal oscillatory potential abnormalities in diabetes and early diabetic retinopathy. *Invest Ophthal Vis Sci* 2004; 45 : 3259-3265.
- [8] Hove MN, Kristensen JK, Lauritzen T, Bek T. The prevalence of retinopathy in an unselected population of type 2 diabetes patients from Arhus County, Denmark. *Acta Ophthalmol Scand* 2004; 82: 443-448
- [9] Huang C, Kim Y, Caramori ML. Cellular basis of diabetic nephropathy: II. The transforming growth factor-beta system and diabetic nephropathy lesions in type 1 diabetes. *Diabetes* 2002; 51: 3577-3581.
- [10] Shukla N, Angelini GD, Jeremy J. Homocysteine as a risk factor for nephropathy and retinopathy in type 2 diabetes. *Diabetologia* 2003; 46: 766-772.
- [11] Svensson M, Eriksson JW, Dahlquist G. Early glycemic control, age at onset, and development of microvascular complications in childhood-onset type 1 diabetes: a population-based study in northern Sweden. *Diabetes Care* 2004; 27: 955-962.
- [12] Saeely CH, Aczel S, Marte T. Cardiovascular complications in type 2 diabetes mellitus depend on the coronary angiographic state rather than on the diabetes state. *Diabetologia* 2004; 47: 145-146.
- [13] Wallace C, Reiber GE, LeMaster J. Incidence of falls, risk factors for falls, and fall-related fractures in individuals with diabetes and a prior foot ulcer. *Diabetes Care* 2002; 25: 1983-1986.
- [14] Centers for Disease Control and Prevention (CDCP). History of foot ulcer among persons with diabetes---United States, 2000-2002. *Morbidity & Mortality Weekly Report*. 2003; 52: 1098-1102.
- [15] World Health Organization Expert Committee on Diabetes Mellitus. Second WHO Technical Report, Series 310. Geneva: World Health Organization, 1965.
- [16] National Diabetes Data Group. Classification and diagnosis of diabetes mellitus and other categories of glucose intolerance. *Diabetes* 1979; 18: 1039-1057.
- [17] Oberley LW. Free radicals and diabetes. *Free Radic Biol Med*. 1988;5(2):113-24.
- [18] Bashan N, Kovsan J, Kachko I, Ovadia H, Rudich A. Positive and negative regulation of insulin signaling by reactive oxygen and nitrogen species. *Physiol Rev*. 2009; 89 (1):27-71.
- [19] Zimmet P, Dowse G, Finch C, Serjeantson S, King H. The epidemiology and natural history of NIDDM lessons from the South Pacific. *Diabetes Metab Rev* 1990; 6: 91-124.
- [20] Eriksson J, Saloranta C, Widen E, et al. Non-esterified fatty acids do not contribute to insulin resistance in persons at increased risk of developing type 2 (noninsulin-dependent) diabetes mellitus. *Diabetologia*. 1991; 34:192-197.
- [21] World Health Organization Study Group. Diabetes Mellitus: WHO Technical Report, Series 727. Geneva: World Health Organization, 1985.
- [22] Zimmet P, Cowie C, Ekoe JM, Shaw JE. Classification of diabetes mellitus and other categories of glucose intolerance. In: *International Textbook of Diabetes Mellitus* chapter 1, 3rd Ed, 2004, pp 3-14
- [23] Atkinson MA, Maclaren NK. The pathogenesis of insulin-dependent diabetes mellitus. *N Engl J Med* 1994; 331: 1428-1436.
- [24] Betterle C, Zannette F, Pedini B. Clinical and subclinical organ-specific autoimmune manifestations in type 1 (insulin-dependent) diabetic patients and their first-degree relatives. *Diabetologia* 1983; 26: 431-436
- [25] Amos A, McCarty D, Zimmet P. The rising global burden of diabetes and its complications, estimates and projections to the year 2010. *Diabetic Med* 1997; 14: S1-S85.
- [26] King H, Aubert R, Herman W. Global burden of diabetes, 1995-2025. Prevalence, numerical estimates and projections. *Diabetes Care* 1998; 21: 1414-1431.
- [27] Harris MI, Flegal KM, Cowie CC, Eberhardt MS, Goldstein DE, Little RR, Wiedmeyer HM, Byrd-Holt DD. Prevalence of diabetes, impaired fasting glucose, and impaired glucose tolerance in U.S. adults: the third National Health and Nutritional Examination Survey, 1988-1994. *Diabetes Care*, 1998; 21: 518-524.
- [28] Report of World Health Organization Study Group. WHO Technical Report Series 1994; World Health Organization, 844: Geneva, Prevention of Diabetes Mellitus.
- [29] Zimmet P. Type 2 (non-insulin-dependent) diabetes: an epidemiological overview. *Diabetologia* 1982; 22: 399-411.
- [30] Christacopoulos PD, Karamanos BG, Tountas CD. The prevalence of diabetes mellitus and non-diabetic glycosuria in a rural population in Greece. *Diabetologia* 1975; 11: A-335.
- [31] DeFronzo RA, Bonadonna RC, Ferrannini E. Pathogenesis of NIDDM. In Albert KGMM, Zimmet P, DeFronzo RA (eds) *International Textbook of Diabetes Mellitus*, 2nd edn. Chichester: Wiley, 1997, pp 635-712.
- [32] Zimmet P, Alberti KGMM, Shaw T. Global and Social implications of diabetes epidemic. *Nature* 2001; 414: 782-787.
- [33] Miller CD, Phillips LS, Ziemer DC, et al. Hypoglycaemia in patients with type 2 diabetes mellitus. *Arch Intern Med*; 2001; 161: 1653-1659.
- [34] American Diabetes Association. Gestational diabetes mellitus. *Diabetes Care* 2001; 24 (suppl 1): S77-S79.
- [35] Raffel LJ, Scheuner MT, Rotter JI. Genetics diabetes. In: Porte D Jr, Sherwin RS, eds. *Ellenberg & Rifkin's Diabetes Mellitus*, 5th ed. Stamford, CT, Appleton & Lange, 1997; 401-454.
- [36] American Diabetes Association. Gestational diabetes mellitus. *Diabetes Care* 2001; 24 (suppl 1): S5-S19.
- [37] Pandit Mk, Burke J, Gustafson AB. Drug-induced disorders of glucose tolerance. *Ann Intern Med* 1993; 118:529-539.
- [38] Olokoba AB, Obateru OA, Olokoba LB, Type 2 Diabetes Mellitus: A Review of Current Trends, *Oman Med J*, (2012); 27(4): 269-273

- [39] Zimmet P, Alberti, KG. Global and societal implications of the diabetes epidemic, Shaw J Nature, (2001); 414(6865):782-787.
- [40] Wild S, Roglic, G, Green A, Sicree R, King H. Global prevalence of diabetes: estimate for the year 2000 and projections for 2030. Diabetes Care, (2004);127(5):1047-1053
- [41] Ramchandran R, Das AK, Joshi SR. Current status of diabetes in India and need for novel therapeutic agents, Suppliment to Japi, (2010);58: 7-9.
- [42] Joshi SR, Parikh RM. India - diabetes capital of the world: now heading towards hypertension. J Assoc Physicians India. (2007); 55: 323-4
- [43] Kumar A, Goel MK, Jain RB, Khanna P, Chaudhary V. India towards diabetes control: Key issues. Australas Med J, (2013); 6(10): 524-31
- [44] Dalle-Donne I, Rossi R, Colombo R, Giustarini D and Milzani A. Biomarkers of oxidative damage in human disease. Clinical Chemistry 2006;52 (4): 601.
- [45] Halliwell B (2001) Role of free radicals in the neuro degenerative diseases: therapeutic implications for antioxidant treatment. Drugs and Aging 18 (9): 685-716.
- [46] Halliwell B and Gutteridge JMC. Free Radicals in Biology and Medicine, 5th edn. Oxford, UK: Oxford University Press. (2015).
- [47] Phaniendra.A, Dinesh Babu Jestadi D B, Periyasamy L, Free Radicals: Properties, Sources, Targets, and Their Implication in Various Diseases. Ind J Clin Biochem (Jan-Mar 2015) 30(1):11-26; DOI 10.1007/s12291-014-0446-0
- [48] Finkel T, Holbrook NJ. Oxidants, oxidative stress and the biology of ageing. Nature. 2000;408:239-47.
- [49] Starkov AA. The role of mitochondria in reactive oxygen species metabolism and signaling. Ann NY Acad Sci. 2008;1147:37-52
- [50] De Duve C, Bauduvin P. peroxisomes (microbodies and related particles). Physiol Rev. 1966;46:323-57.
- [51] Schrader M, Fahimi HD. Review Peroxisomes and oxidative stress. Biochim Biophys Acta. 2006;1763(12):1755-66.
- [52] Cheeseman KH, Slater TF. An introduction to free radical biochemistry. Br Med Bull. 1993;49(3):481-93
- [53] McCord JM, Fridovich I. The reduction of cytochrome c by hypoxanthine and xanthine oxidase. Biochem. J. 1968, 243, 5753-5760.
- [54] Mccord JM, Fridovich I. Superoxide dismutase: An enzymic function for erythrocuprein (hemocuprein). J. Biol. Chem. 1969, 244, 6049-6055.
- [55] Boveris BA, Oshino N, Chance B. The cellular production of hydrogen peroxide. Biochem. J. 1972, 128, 617-630
- [56] Halliwell B. Free radicals, reactive oxygen species and human disease: A critical evaluation with special reference to atherosclerosis. Br. J. Exp. Pathol. 1989, 70, 737-757.
- [57] Schwab L, Goroncy. L, Palaniyandi S, Gautam S, Triantafyllou A, Mocsaí A, Reichardt W, Karlsson FJ, Radhakrishnan SV, Hanke K. Neutrophil granulocytes recruited upon translocation of intestinal bacteria enhance graft-versus-host disease via tissue damage. Nat. Med. 2014; 20, 648-654
- [58] Azizova OA, Panasenka O, Volnov T, Vladimirov YA. Free radical lipid oxidation affects cholesterol transfer between lipoproteins and erythrocytes. Free Radic. Biol. Med. 1989; 7, 251-257.
- [59] Lyras L, Perry RH, Perry EK, Ince PG, Jenner P, Jenner P, Halliwell B. Oxidative damage to proteins, lipids, and DNA in cortical brain regions from patients with dementia with Lewy bodies. J. Neurochem. 1998; 71, 302-312.
- [60] Halliwell B, Aruoma OI. DNA damage by oxygen-derived species. FEBS Lett. 1991; 281, 9-19.
- [61] Cadet J, Douki T, Ravanat JL. Oxidatively generated base damage to cellular DNA. Free Radic. Biol. Med. 2010; 49, 9-21.
- [62] Sies H. Oxidative stress: Introductory remarks. In Oxidative Stress; Academic Press: London, UK, 1985; pp. 1-8.
- [63] Rosen H, Klebanoff SJ, Wang Y, Brot N, Heinecke JW, Fu X. Methionine oxidation contributes to bacterial killing by the myeloperoxidase system of neutrophils. Proc. Natl. Acad. Sci. USA 2009; 106, 18686-18691.
- [64] Cadet J, Wagner JR. Oxidatively generated base damage to cellular DNA by hydroxyl radical and one-electron oxidants: Similarities and differences. Arch. Biochem. Biophys. 2014; 557, 47-54.
- [65] Salgo MG, Squadrito GL, Pryor WA. Peroxynitrite causes apoptosis in rat thymocytes. Biochem. Biophys. Res. Commun. 1995; 215, 1111-1118.
- [66] Panasenka OM, Evgina SA, Driomina ES, Sharov VS, Sergienko VI, Vladimirov YA. Hypochlorite induces lipoproteins and lipid peroxidation in blood phospholipid liposomes. Free Radic. Biol. Med. 1995; 19, 133-140
- [67] Chevion M, A site-specific mechanism for free radical induced biological damage: The essential role of redox-active transition metals. Free Radic. Biol. Med. 1988; 5, 27-37.
- [68] Violi F, Milite MT, Medica IC, La A. Nitric oxide and its role in lipid peroxidation. Diabetes Metab. Res. Rev. 1999; 15, 283-288.
- [69] Miller DM, Buettner GR, Aust SD. Transition metals as catalysts of "autoxidation" reactions. Free rad. Biol. 1990; 8, 95-108
- [70] Cadenas E, Sies H. The lag phase. Free rad. 1998; 601-609
- [71] Valko M, Morris H, Cronin MTD. Metals, toxicity and oxidative stress. Curr. Med. Chem. 2005;12, 1161-1208.
- [72] Muller FI, Liu Y, Van remmen H. Complex III releases superoxide to both sides of the inner mitochondrial membrane. J. boil. Chem., 279, 49064-49073.
- [73] Liochev S.I., Fridovich I. (2002), The Haber-Weiss cycle-70 year later. An alternative view. Redox, 2004; 55-57.
- [74] Pastor N, Weinstein H, Jamison E, Brenowitz M. A detailed interpretation of OH radical footprints in a TBP DNA complex reveals the role of dynamics in the mechanism of sequence-specific binding. J. Mol. Biol. 2000; 304, 55-68.
- [75] Rae TD, Schmidt PJ, Pufahl RA, Halloran TV. Undetectable intracellular free copper. The requirement of a copper chaperone for superoxide dismutase. Science. 1999; 284, 805-808
- [76] Kakhlon O, Cabantchik ZI. the labile iron pool: Characterization, measurement, and participation in cellular processes. Free Radic. Biol. Med. 2002;33, 1037-1046
- [77] De Grey ANDJ, The forgotten radical. DNA Cell. Boil. 2002; 21, 251-257
- [78] Aikens J, Dix TA, Peroxyl radical (HOO*) initiated lipid peroxidation- The role of fatty hydroperoxides. J. Biol. Chem, 1991; 266, 15091-15098
- [79] Bergendi L, Benes L, Durackova Z, Ferencik M. Chemistry, physiology and pathology of free radicals. Life, sci, 1999; 65, 1865-1874
- [80] Chiu CC. Neuroprotective properties of nitric oxide, Ann. N.Y. Acad sci, 1999; 890, 301-311
- [81] Klatt P, Lamas S. Regulation of protein function by S-glutathiolation in response to oxidative and nitrosative stress. Eur. J. Biochem, 2000; 267, 4928-4944



International Journal of Recent Development in Engineering and Technology
Website: www.ijrdet.com (ISSN 2347-6435 (Online) Volume 15, Issue 08, August 2026)

- [82] Ridnour LA, Thomas DD, Mancardi D, Espey MG, Miranda KM, Paolucci N. The chemistry of nitrosative stress induce by nitric oxide and reactive nitrogen oxide species. Putting perspective on stressful biological situation. *Biol. Chem*, 2004; 385, 1-10.
- [83] Carr A, McCall MR, Frei B. Oxidation of LDL by myeloperoxidase and reactive nitrogen species reaction pathways and antioxidant protection. *Arterioscl. Thromb vasc. Biol.* 2000; 20, 1716-1723.
- [84] Donath MY, Størling J, Maedler K, Mandrup-Poulsen T. Inflammatory mediators and islet beta-cell failure: A link between type 1 and type 2 diabetes. *J Mol Med (Berl)* 2003;81(8):455-70.
- [85] Pinkse GG, Tysma OH, Bergen CA, Kester MG, Ossendorp F, van Veelen PA. Autoreactive CD8 T cells associated with beta cell destruction in type 1 diabetes. *Proc Natl Acad Sci U S A* 2005;102:18425-30.
- [86] Pickup JC. Inflammation and activated innate immunity in the pathogenesis of type 2 diabetes. *Diabetes care* 2004;27:813-23.
- [87] Wiley J. Evaluation of some biochemical changes in diabetic patients. *Diabetes Care* 1992;15(10):413-22.
- [88] Wellen KE, Hotamisligil GS. Obesity-induced inflammatory changes in adipose tissue. *J Clin Invest* 2003;112(12):1785-8.
- [89] Akkus I, Kalak S, Vural H, Caglayan O, Menekse E, Can G. Leukocyte lipid peroxidation, superoxide dismutase, glutathione peroxidase and serum and leukocyte vitamin C levels of patients with type II diabetes mellitus. *Clin Chim Acta* 1996;244(2):221-7.
- [90] Kalaivanam KN, Dharmalingam M, Marcus SR. Lipid peroxidation in type 2 diabetes. *Int J Diabetes Dev Ctries* 2006;26(1):30-2.
- [91] Kashiwagi A. Complications of diabetes mellitus and oxidative stress. *JMAJ* 2001;44(12):521-8.