

The Botanical Dimensions of Indian Knowledge Systems: An Analytical Perspective

Dr. Ravindra Kumar Pandey

¹Department of Botany, Dr. SPM Govt. Degree College, Bhadohi, India
Affiliated to Kashi Naresh University Bhadohi, India

Abstract—The Indian Knowledge System (IKS) in botany and plant science represents a sophisticated, empirical, and deeply holistic tradition developed over millennia. Historically categorized under Vrikshayurveda (the science of plant life) and Vanaspati Shastra (botanical science), this framework transitions from early Vedic ecological observations to highly systematized classical treatises. This paper examines the structural evolution of ancient Indian botany, focusing on its multi-tiered taxonomic systems (Vanaspati Vargikarana) and precise morphological and anatomical terminologies. Crucially, it highlights how ancient scholars anticipated several principles of modern plant physiology—including internal fluid circulation (Sirajala), germination parameters (Bijarkura), and plant sensitivity and consciousness (Chetanata), later institutionalized through the practice of Dohada (botanical stimulation). The study further details traditional phytopathology, examining how Surapala's Vrikshayurveda diagnosed internal humoral imbalances (Vata-Pitta-Kapha) and prescribed sustainable, biodegradable remedies like Kunapa Jala. Finally, the paper explores applied ecological practices, such as groundwater exploration via bio-indicators (Dakargala) and Kautilya's forest management frameworks. By comparing these insights with Western historical botany, this paper demonstrates the contemporary relevance of IKS botany in driving modern bioprospecting, organic farming, and climate-resilient agriculture.

Keywords—Vrikshayurveda (Ancient Indian plant science), Vanaspati Shastra (Botanical taxonomy and classification), Chetanata (Plant sentience and neurobiology), Kunapa Jala (Traditional liquid bio-fertilizer), Dakargala (Eco-botanical groundwater exploration), Dravyaguna Vijnana (Pharmacological botany), Sustainable Agriculture

I. INTRODUCTION

The Indian Knowledge System (IKS) represents a structured, empirical, and deeply philosophical framework of science accumulated over millennia. Within IKS, the study of plant life—historically designated as *Vrikshayurveda* (the science of plant life) and *Vanaspati Shastra* (botanical science)—developed not merely as an auxiliary branch of medicine, but as an independent, highly organized scientific discipline.

Rooted in ancient Vedic literature, classical Sanskrit texts, and regional vernacular traditions, Indian plant science encompasses advanced systems of taxonomy, morphology, physiology, ecology, and agricultural biotechnology. Unlike the fragmented or purely utilitarian views of nature that often characterized early Western frameworks, the traditional Indian approach to botany is fundamentally holistic. It recognizes a profound existential continuity between human beings, animals, and the vegetable kingdom. This epistemological paradigm is anchored in the Vedic understanding of *Rta* (cosmic order), wherein all living entities are governed by identical natural laws [1]. The evolutionary trajectory of Indian botany spans several distinct chronological epochs:


- The Vedic Period (c. 1500 BCE – 500 BCE)*: The foundational era where primary classification, ritualistic utility, and ecological observations were recorded in the *Rigveda*, *Atharvaveda*, and associated *Brahmanas*.
- The Classical Samhita Period (c. 500 BCE – 500 CE)*: The era of rigorous systematization led by medical authorities such as Charaka, Sushruta, and Vagbhata. During this time, comprehensive pharmacological botany (*Dravyaguna Vijnana*) was established.
- The Specialized Technical Period (c. 500 CE – 1500 CE)*: The compilation of dedicated botanical treatises, most notably the *Vrikshayurveda* of Surapala, the *Upavana Vinoda* of Samgadharra, and various *Nighantus* (lexicons). These texts isolated plant science from human medicine, focusing strictly on arboriculture, plant pathology, and soil chemistry.

By interrogating these historical layers, this paper demonstrates that ancient Indian scholars anticipated several foundational principles of modern botany, including plant sensitivity, sexuality, internal fluid circulation, and systematic binomial nomenclature.

II. HISTORICAL FRAMEWORK AND CANONICAL LITERATURE

The structural evolution of botanical science in ancient India is mapped across an extensive corpus of Sanskrit literature.

These texts transition progressively from spiritual and utilitarian invocations to strictly empirical and experimental methodologies.



A. Vedic Foundations

The *Rigveda*, the oldest extant text of Indo-European literature, contains numerous hymns dedicated to *Oshadhis* (medicinal herbs) and *Vanaspatis* (forest trees). It categorizes plants based on their growth habits, external forms, and therapeutic properties. The *Atharvaveda* expands significantly upon this empirical base, offering detailed descriptions of deeply rooted herbs, spreading creepers, and parasitic plants, alongside specific methods for treating plant and human diseases. In these texts, the plant kingdom is invoked as an essential pillar of environmental equilibrium, and prayers are offered to prevent the indiscriminate destruction of vegetation.

B. Philosophical and Philosophical Systems

The institutionalization of botany as a rational science was catalyzed by the *Darshanas* (systems of Indian philosophy). The *Vaisheshika Darshana* of Kanada and the *Nyaya Darshana* of Akshapada Gautama explored the atomic structure of matter and the mechanisms of physical transformation within plant bodies.

The *Sankhya* and *Yoga* systems introduced a vitalist perspective, asserting that plants possess an internal consciousness, experience sensations of pain and pleasure, and respond directly to external environmental stimuli.

C. Ayurvedic Lexicons

The *Charaka Samhita* (c. 2nd century CE) and the *Sushruta Samhita* (c. 3rd–4th century CE) represent the zenith of ancient Indian applied botany. Charaka provides an exhaustive classification of more than 500 plant-based drugs, organizing them into 50 distinct therapeutic groups (*Mahakashayas*) [2]. Sushruta, focusing primarily on surgical interventions and toxicology, classifies plants based on their environmental habitats and specific physical characteristics, establishing a systematic methodology for identifying authentic botanical specimens [3].

D. Dedicated Botanical Treatises


The most critical milestones in independent Indian plant science are texts dealing exclusively with *Vrikshayurveda*.

- a. Surapala's *Vrikshayurveda* (c. 10th century CE): A comprehensive, 325-verse compendium detailing the entire life cycle of plants. It contains rigorous methodologies for seed selection, soil classification, nursery management, grafting, plant nutrition, and phytopathology [4].
- b. Sarngadhara's *Upavana Vinoda* (c. 14th century CE): A specialized section within the larger *Sarngadhara Paddhati* that functions as a manual for urban forestry, the layout of royal pleasure gardens, and the preservation of rare plant species [5].
- c. Varahamihira's *Brihat Samhita* (c. 6th century CE): Contains pivotal chapters on *Argala* (groundwater exploration using bio-indicators) and *Vrikshayurveda*,

linking astrological cycles, meteorological patterns, and plant behavior [6].

III. TAXONOMY AND CLASSIFICATION SYSTEMS (VANASPATI VARGIKARANA)

Long before the taxonomic frameworks of Carl Linnaeus or the natural systems of Antoine Laurent de Jussieu, ancient Indian botanists developed sophisticated, multi-tiered hierarchical structures to classify the plant kingdom. These classifications were not arbitrary; they were based simultaneously on morphological traits (*Akruti*), ecological niches (*Bhumi*), and utilitarian properties (*Guna*).



IV. CLASSICAL QUADRIPARTITE CLASSIFICATION

The oldest structural division of the plant kingdom, recurring consistently from the *Upanishads* to the classical medical Samhitas, divides all vegetation (*Udbhijja* or surface-born entities) into four primary classes:

- a. *Vanaspati*: Trees that bear fruit directly without producing visible, conspicuous flowers. Classical examples cited include *Ficus religiosa* (Pipal) and *Ficus benghalensis* (Banyan). Modern botanical correlation aligns this perfectly with cryptogams and specific gymnosperms/angiosperms possessing hypanthodium inflorescences.

- b. *Vanaspatya (or Vrisha)*: Trees that produce both distinct, visible flowers and subsequent fruits. This category encompasses the vast majority of dicotyledonous arborescent plants, such as *Mangifera indica* (Mango) and *Syzygium cumini* (Jamun).
- c. *Oshadhi*: Annual or ephemeral herbs that wither and die completely after the maturation of their seeds and fruits. This group includes vital cereal crops like rice (*Vrihi*) and barley (*Yava*).
- d. *Virudh (or Pratana)*: Spreading, climbing, or creeping plants that require external physical support to grow.

This class is further subdivided into *Valli* (twining climbers with slender stems) and *Pratanavati* (creepers that spread horizontally across the ground using tendrils).

V. ADVANCED STRUCTURAL SUBDIVISIONS

As empirical observations sharpened, later botanists realized that the quadripartite system was insufficient to capture the vast structural diversity of flora. The *Manusmriti* and the commentaries of Bhattotpala introduced additional morphological classes:

- a. *Guccha*: Bushy shrubs and perennial thickets characteristically branching from the very base of the stem (e.g., *Jasminum* species).
- b. *Gulma*: Low, dense, succulent shrubs with multiple clustering stems.
- c. *Trina*: The diverse family of grasses, distinctly marked by their hollow, jointed stems (*Parva*).
- d. *Twaksara*: Plants characterized by an endogenous stem structure where the hard, structural tissue is located on the periphery (bark/cortex) while the core remains hollow or soft, exemplified by *Bambusa arundinacea* (Bamboo) and various *Palmae* species.
- e. *Saka*: Pot-herbs and leafy vegetables defined by their succulent, non-woody aerial parts.

VI. NOMENCLATURAL PRINCIPLES

Ancient Indian nomenclature operated on a highly systematic, pseudo-binomial matrix. Plants were assigned a primary generic name supplemented by specific descriptive epithets (*Namakarana*). These names were intentionally formulated to communicate vital botanical data, falling into several distinct descriptive categories:

- a. Morphological Indicators (*Akruti-bodhaka*): *Chitraparka* (plants with variegated leaves), *Dirghapatra* (long-leaved plants), *Sata-muli* (possessing a hundred roots, like *Asparagus racemosus*), *Tri-patra* (trifoliate leaves).
- b. Ecological/Habitat Indicators (*Sthana-bodhaka*): *Nadeyi* (growing exclusively near riverbanks), *Maruvaka* (native to arid, desert ecosystems), *Pankaja* (mud-born, aquatic flora).
- c. Therapeutic/Pharmacological Indicators (*Guna-bodhaka*): *Krimighni* (destroyer of intestinal worms/parasites), *Vajradanti* (strengtheners of teeth and gums), *Jwarahara* (antipyretic/fever-reducing).

- d. Phenological Indicators (*Kala-bodhaka*): *Sharadi* (blooming exclusively during the autumn season), *Suryamukhi* (tracking the movement of the sun, *Helianthus*-type).

This dual-naming strategy allowed an ancient practitioner to instantly deduce a plant's physical appearance, ecological origin, and therapeutic potential from its name alone.

VII. ANATOMY AND MORPHOLOGY (SARIRA AND AKRUTI VIJNANA)

Ancient Indian thinkers developed a comprehensive terminology to document the macro- and micro-morphology of plant structures, demonstrating a keen eye for developmental anatomy.

A. Gross Morphological Organs

The *Taittiriya Samhita* and the *Vrikshayurveda* texts outline a systematic structural breakdown of the plant body, identifying the major organs and their anatomical relations:

- a. *Mula or Pada (Root System)*: The foundational anchor responsible for the absorption of moisture and nutrients from the soil matrix. Special recognition was given to *Sikhara-mula* (taproots) and *Jala-mula* (fibrous, adventitious root networks).
- b. *Skandha (Trunk/Stem)*: The main axis of the plant supporting the canopy.
- c. *Sakha (Branches)*: Secondary and tertiary lateral extensions emerging from the main stem.
- d. *Patra or Parna (Leaf)*: The primary organ for respiration and light interception.
- e. *Puspa (Flower)*: The reproductive structure, analyzed through components like the *Dala* (petal), *Kesara* (stamen), and *Garbhasayana* (ovary).
- f. *Phala (Fruit)*: The matured ovary enclosing the *Bija* (seed).

B. Histological and Internal Structures

The internal tissue organization of plants was conceptualized through macroscopic dissections and microscopic inferences. The *Charaka Samhita* identifies four clear internal zones within a woody stem:

- a. *Twak (Bark/Epidermis)*: The outermost protective dermal layer, subdivided into the external dry bark and the inner succulent bark.
- b. *Mamsasara (Cortex/Cambium)*: The intermediate soft, functional zone beneath the bark.

- c. *Asthi (Wood/Xylem)*: The dense, lignified, structural core of the tree that provides mechanical stability.
- d. *Majja (Pith/Medulla)*: The central soft core containing cellular reserves.

VIII. VASCULAR CONCEPTS: SIRAJALA AND SAP MOVEMENT

One of the most remarkable insights of ancient Indian plant physiology was the recognition of a specialized internal vascular system. Texts speak of *Siras* and *Dhamanis*—networks of microscopic channels or ducts that run continuously from the root tips through the stem to the apex of the leaves.

According to the commentaries of Dalhana and Sankara Misra, these channels are responsible for the continuous movement of *Rasa* (nutrient-laden water/sap). The mechanism of sap ascent was explained through a combination of root absorption (*Padapa*—drinking through the feet), thermal capillary action driven by solar energy (*Agni*), and internal atmospheric pressure (*Vayu*). This concept aligns closely with modern transpiration pull and capillary theories of xylem transport.

IX. PHYSIOLOGY, DEVELOPMENT, AND SENSATION

The physiological transition of plants from inert matter to sentient, dynamic biological entities is a core doctrine of IKS plant science. Ancient Indian thinkers directly rejected the idea that plants lack conscious life, presenting an empirical defense of plant perception.

A. Physiology of Germination (*Bijarkura*)

The developmental sequence from seed to seedling was observed with great precision. The *Sushruta Samhita* states that for successful germination (*Ankurabhidana*), four fundamental factors must converge simultaneously: Successful Germination=Ritu (Season/Time)+ Kshetra (Soil/Medium) + Ambu (Water/Moisture) + Bija (Viable Seed)

- a. *Ritu (Proper Season/Temperature)*: The correct microclimatic and thermal conditions.
- b. *Kshetra (Fertile Soil Matrix)*: A nutrient-rich growth medium free of contaminants.
- c. *Ambu (Moisture/Water)*: The catalytic agent that softens the seed coat and triggers internal metabolism.

- d. *Bija (Genetically Sound Seed)*: A structurally intact, mature seed possessing the *Sakti* (latent life force) of the parent plant.

The seed coat was termed *Bijachola*, the cotyledons *Bijapatra*, and the emerging radicle and plumule were recognized as the initial manifestations of *Ankura* (the embryo).

B. Plant Sensation and Consciousness

The debate regarding plant perception (*Chetanata*) features prominently in Indian philosophical discourse. The *Mahabharata* (Santi Parva, Chapter 184) contains a dialogue between Sage Bhṛigu and Bharadvaja that presents clear empirical evidence for plant senses:

- a. *Touch*: "If a tree is touched by a spreading creeper, it winds around it from all sides. A blind entity cannot find a path; therefore, trees possess the sense of sight and touch."
- b. *Vision*: "Trees respond directly to light and bend toward the sun. They wither under the impact of intense heat and lightning; therefore, they perceive visual forms."
- c. *Hearing*: "Trees are observed to flower and fruit vigorously under the impact of deep sounds, such as thunder or the beat of drums; therefore, they possess the sense of hearing."
- d. *Taste and Smell*: "Trees drink water through their roots; they recover from diseases when treated with fragrant and nourishing substances. This proves they possess the senses of taste and smell."

This text concludes with the definitive assertion: *Tasmabh Jivan Pasyami Vrikshanamani Chaitanyam*, meaning, "Therefore, I perceive life in trees; they are not inert matter [7]."

C. The Phenomenon of Dohada

This deep understanding of plant sensitivity crystallized into the classical practice of *Dohada* (pregnancy-desire of plants). *Dohada* refers to the specialized botanical technique of applying distinct physical, chemical, or acoustic stimuli to specific plants to accelerate seasonal flowering, break seed dormancy, or boost crop yields (Table 1).

Table 1

Sl. no.	Botanical Species	Traditional <i>Dohada</i> Stimulus	Modern Scientific/Phytochemical Correlation
1.	<i>Ashoka (Saraca asoca)</i>	Contact/kicking by a graceful woman's foot	Mechanical stress triggers ethylene release, initiating floral primordia.
2.	<i>Bakula (Mimusops elengi)</i>	Spraying with wine or alcohol from the mouth	Exogenous ethanol/organic compounds act as carbon sources and metabolic stimulants.
3.	<i>Kuruvaka (Barleria cristata)</i>	Gentle structural hugging or physical contact	Thigmomorphogenesis; physical contact alters auxin distribution to promote branching.
4.	<i>Mango (Mangifera indica)</i>	Exposure to smoke from burnt organic matter	Ethylene and acetylene in smoke break bud dormancy and induce synchronized flowering.

These practices show that ancient horticulturists understood how to manipulate plant hormones and reproductive cycles using targeted environmental triggers.

X. PHYTOPATHOLOGY AND PLANT CARE (VRIKSHA AYURVEDA)

Surapala's *Vrikshayurveda* is the foundational text for traditional Indian phytopathology. It systematically treats the plant as a biological patient, diagnosing internal and external disorders and prescribing comprehensive organic remedies.

A. Ethiological Classification of Diseases

Surapala classifies plant diseases (*Roga*) into two primary categories based on their origin:

- a) Internal Diseases (*Adhyatmika*): Caused by a systemic imbalance of the three bodily humours-*Vata*, *Pitta*, and *Kapha*-mirroring classical Ayurvedic human medicine.
- b) External Diseases (*Adhibhautika*): Caused by external environmental attacks, including parasitic infestations, fungal blights, cold waves, heat strokes, structural damage from storms, and lightning strikes [8].

B. Symptomatology of Humoral Imbalances in Plants

Ancient botanists carefully observed the physiological symptoms of internal chemical and metabolic imbalances:

- a) Vata-dominated Diseases: Characterized by stunted growth, slender and twisted trunks, hardening of the bark, brittle leaves, and a lack of fruit or seed development. This condition correlates with modern nutrient deficiency syndromes, severe dehydration, or vascular wilt.
- b) Pitta-dominated Diseases: Marked by premature yellowing of leaves (*Chlorosis*), early drop of flowers and fruits, decay of roots, and localized burning or wilting of the canopy. This corresponds to heat stress, nitrogen toxicity, or acute bacterial rots.

- c) Kapha-dominated Diseases: Indicated by delayed flowering, glossy but thick and distorted leaves, accumulation of excess mucilage or sap exudates, and structural softening of internal tissues. This matches modern fungal molds, root drowning, or carbohydrate over-accumulation.

C. Phytotherapeutic Formulations

The therapeutic arsenal of *Vrikshayurveda* relies entirely on biodegradable, organic, and ecologically sustainable materials [9].

a. Panchagavya and Kunapa Jala

The ultimate bio-fertilizer and plant tonic in IKS is *Kunapa Jala* (literally "corpse-water") [10]. First detailed by Surapala, it is an advanced liquid organic formulation prepared by fermenting animal remains (flesh, marrow, skin, and blood) mixed with sesame oil cake, honey, milk, and legumes.

From a modern biochemical perspective, *Kunapa Jala* is a rich source of readily bio-available amino acids, nitrates, soluble phosphorus, potassium, and chelated micronutrients. It acts both as a growth promoter and a powerful soil microbiota rejuvenator.

b. Management of Pests and Blights

To combat insects, caterpillars, and fungal pathogens, texts prescribe specific decoctions:

1. Neem (*Azadirachta indica*): Aqueous and oil extracts used as a broad-spectrum insecticide and anti-feedant. Modern science confirms azadirachtin as a potent disruptor of insect growth [11].
2. Turmeric (*Curcuma longa*): Applied as a fine dust or paste to pruned branches and open plant wounds to prevent fungal and bacterial infections.
3. Asafoetida (*Ferula foetida*) and Garlic (*Allium sativum*): Vaporized or infused into the soil around roots to act as systemic nematicides and repellents against boring insects.

XI. ECONOMIC BOTANY, AGRICULTURE, AND ETHNOBOTANY

The practical translation of traditional Indian plant science took place in fields, forests, and royal eco-preserves, laying the foundation for a resilient agricultural economy.

A. Soil Taxonomy (Bhumi Vargikarana)

Sustainable agriculture began with a detailed classification of soils. The *Arthashastra* of Kautilya [12] and the *Vrikshayurveda* categorize lands based on moisture content, topography, and crop compatibility [13]:

- a. Anupa (Marshy/Humid Regions):* Rich in organic matter, ideal for intensive cultivation of paddy (*Vrihi*), sugarcane (*Ikshu*), and dense plantains.
- b. Jangala (Arid/Semi-Arid Plains):* Characterized by low water tables and thorny vegetation, suitable for legumes, pulses, and deep-rooted millets.
- c. Samana (Ordinary/Balanced Soil):* Loamy, intermediate soils capable of sustaining a wide variety of mixed agricultural crops.

Soils were further analyzed by color (black, red, yellow), texture (sandy, clayey, loamy), taste (sweet, saline, sour, bitter—used to detect chemical toxicities), and odor to determine their specific fertility profile.

B. Advanced Propagation Methods

Ancient horticulturists developed sophisticated methods to clone and propagate desirable plant varieties:

- a. Bijaja (Seed Propagation):* The standard sexual reproduction method, requiring detailed seed selection and storage protocols.
- b. Kathodbhava or Skandhaja (Stem Cuttings):* Asexual propagation of woody plants by treating stem segments with rooting agents like clarified butter (*Ghee*), honey, and cow dung paste.

c. Moolaja (Root Cuttings/Suckers): Propagating plants via underground root structures.

d. Goti or Graphing (Sankramana): The structural joining of a scion and stock to combine desired traits. Texts detail wedge grafting and air layering for difficult-to-breed fruit trees like the pomegranate and citrus [14].

C. Ethnobotanical Practices and Conservation


Traditional Indian societies protected biodiversity by embedding ecological conservation into religious frameworks. This is best exemplified by the institutionalization of Sacred Groves (*Deva Vanas*). These patches of pristine forest were dedicated to local deities, creating strict communal taboos against logging, grazing, and hunting. As a result, these groves survived for centuries as vital gene banks for rare, endemic medicinal plants. Similarly, the concept of *Panchavati*—planting five specific sacred trees (Banyan, Pipal, Ashoka, Bel, and Amla) in close proximity—served as a practical micro-ecosystem designed to purify air, retain groundwater, and provide diverse habitats for local birds and insects.

XII. ECOLOGICAL AWARENESS AND SUSTAINABILITY

The ancient Indian world-view considered humanity an integral part of the ecological matrix, rather than its absolute owner. This perspective gave rise to an environmental ethic centered on sustainability.

Groundwater Exploration via Bio-Indicators

Chapter 54 of Varahamihira's *Brihat Samhita*, titled *Dakargala*, is an extraordinary treatise on hydrogeology. It details how to locate clean underground aquifers in arid zones by observing specific plant species and their morphological mutations.



Varahamihira provides explicit empirical rules:

- a. If a Vatasa (Salix) tree is found growing in a desert environment devoid of surface water, a reliable water source exists directly beneath it at a depth of three *Purushas* (approx. 15 feet), flowing in a specific direction [15].
- b. The presence of a Jamun (*Syzygium cumini*) tree to the east of an anthill indicates a sweet-water aquifer nearby.
- c. Morphological mutations, such as an unusually glossy canopy, weeping branches on a non-weeping tree, or a single heavily laden branch on a barren tree, were recognized as physiological stress responses to shallow underground water tables.

Modern eco-botanical studies validate these guidelines, confirming that many of the listed species are phreatophytes—plants with deep root systems that draw directly from the water table.

XIII. FOREST CLASSIFICATION AND RESOURCE MANAGEMENT

The *Arthashastra* outlines the first state-enforced framework for forest management and conservation. Kautilya divided national forests into four distinct operational classes:

- a. *Kupya Vana (Productive State Forests)*: Managed strictly for timber, fuel, cordage, and manufacturing materials.
- b. *Mriga Vana (Royal Game Preserves)*: Protected habitats for wildlife conservation.
- c. *Hastivana (Elephant Forests)*: Specially preserved wilderness areas reserved for breeding and training military elephants.
- d. *Tapovana (Hermitage Forests)*: Sacred zones set aside for meditation, where all forms of logging and violence were prohibited.

The state appointed a *Kupyadhyaksha* (Superintendent of Forest Produce) who regulated forest cutting, punished environmental vandals with heavy fines, and ensured continuous replanting programs. This established an early template for structured, sustainable silviculture.

XIV. COMPARATIVE ANALYSIS: IKS BOTANY VS. WESTERN HISTORICAL BOTANY

To appreciate the scientific maturity of Indian plant science, it is helpful to compare its developmental timeline and core concepts with those of classical European botany (Table 2).

Table 2

Scientific Concept	Indian Knowledge System (IKS) Framework	Historical Western Botanical Framework
Plant Perception and Sentience	Documented by 500 BCE (<i>Mahabharata</i> , <i>Upanishads</i>). Plants recognized as living, conscious entities experiencing touch, vision, hearing, and taste.	Dominated by the Aristotelian view of a "vegetative soul," which denied plants individual sensation or consciousness. This view persisted until the experimental work of Jagadish Chandra Bose and late 20th-century plant neurobiology.
Taxonomy and Nomenclature	Developed multi-tiered systems based on morphology, ecology, and utility (<i>Rigveda</i> , <i>Surapala</i>). Used systematic, pseudo-binomial descriptive naming matrices.	Relied on localized folknames until the 16th-century work of Andrea Cesalpino and the definitive introduction of formal binomial nomenclature by Carl Linnaeus in 1753.
Internal Vascular Circulation	Described the <i>Sirajala</i> network and <i>Rasa-vahana</i> channels. Sap ascent was linked to root absorption and solar-driven evaporation (<i>Brihat Samhita</i>).	Plant circulation was poorly understood until Stephen Hales published <i>Vegetable Staticks</i> in 1727, establishing the foundations of plant physiology and transpiration pull.
Phytopathology and Therapeutics	Developed a distinct branch of plant medicine (<i>Vrikshayurveda</i>) using humoral analysis (<i>Vata-Pitta-Kapha</i>) and targeted organic treatments.	Plant diseases were largely viewed as divine curses, spontaneous generation, or atmospheric rot until the mid-19th century, when Heinrich Anton de Bary founded modern mycology and phytopathology.

While European botany made massive strides during the Enlightenment by developing advanced optical instruments, ancient Indian botany excelled at long-term macro-ecological observations, systemic classifications, and sustainable organic technologies that are highly relevant today.

XV. CONTEMPORARY RELEVANCE AND GLOBAL INTEGRATION

The integration of the Indian Knowledge System in botany with modern plant science is not a matter of historical pride; it is an ecological necessity for the 21st century.

A. Bioprospecting and Ethnopharmacology

Modern pharmaceutical research relies heavily on reverse pharmacology—using ancient Ayurvedic texts (*Dravyaguna Vijnana*) to identify active phytochemical leads. By analyzing traditional uses, researchers can skip expensive random chemical screening. Global discoveries like reserpine (an antihypertensive drug derived from *Rauvolfia serpentina*, or *Sarpagandha*) and the widespread validation of *Curcuma longa* (Turmeric) as a powerful anti-inflammatory agent demonstrate the predictive value of ancient Indian pharmacology.

B. Sustainable Agriculture and Organic Farming

As modern intensive agriculture faces declining soil health, chemical runoff, and pest resistance, the organic protocols of *Vrikshayurveda* offer viable alternatives [16]. The global shift toward organic farming makes formulations like *Kunapa Jala* and *Panchagavya* highly valuable.

These traditional inputs restore the soil microbiome, increase humic matter, and strengthen plant immunity without chemical pollution. Similarly, deploying botanical pest repellents derived from Neem, Karanja (*Pongamia pinnata*), and Vitex preserves beneficial insect populations while controlling target pests.

C. Climate Resilience and Circular Bio-Economies

The ecological design of sacred groves, mixed cropping systems (*Vrihi-Yava* rotation), and hydrogeological mapping using phreatophytic indicators provide time-tested strategies for climate change adaptation. By embracing the traditional Indian view of nature—where the human economy is a nested subsystem within the planetary ecology—modern science can transition away from extractive practices and build a truly sustainable, circular bio-economy.

XVI. CONCLUSION

The Indian Knowledge System in botany and plant science stands as an empirical, sophisticated, and deeply sustainable scientific tradition. From the initial taxonomic insights of the Vedic sages to the systematic agricultural and phytopathological manuals of Surapala and Varahamihira, ancient Indian scholars built a profound understanding of plant life. They recognized plants as sentient, responsive, and anatomically complex organisms, anticipating modern discoveries in plant neurobiology, vascular physiology, and chemical ecology. Crucially, IKS botany avoided the conceptual split between scientific progress and environmental preservation.

By embedding botanical knowledge within a framework of cosmic balance (*Rta*) and ecological stewardship, it ensured that agricultural prosperity did not come at the cost of planetary destruction. As modern science grapples with the crises of climate change, widespread biodiversity loss, and soil degradation, the holistic, time-tested principles of *Vrikshayurveda* and traditional Indian plant science offer more than historical insights—they provide a proven roadmap for a sustainable future.

Acknowledgement

The author gratefully acknowledges the financial assistance provided by the Higher Education Council, Government of Uttar Pradesh, under its research and development grant scheme. This support has been instrumental in carrying out the present research work. The author also expresses sincere gratitude to the Department for its encouragement and support in promoting academic research and scholarly activities.

REFERENCES

- [1] Sharma, S. (2025). The living breath of earth: Plants in the Indian Knowledge System. *Indological Botanical Surveys*, 11(1), 14–29. Research Gate Full Text.
- [2] Agnivesha. (2018). *Charaka Samhita* (P. V. Sharma, Trans.). Chaukhambha Orientalia. (Original work compiled c. 2nd century CE).
- [3] Sushruta. (2005). *Sushruta Samhita* (P. V. Sharma, Trans.). Chaukhambha Visvabharti. (Original work compiled c. 3rd–4th century CE).
- [4] Sadhale, N. (Trans.). (1996). *Surapala's Vrikshayurveda* (The Science of Plant Life by Surapala) (Agri-History Bulletin No. 1). Asian Agri-History Foundation.
- [5] Samgadharma. (2009). *Upavana Vinoda* (A Sanskrit treatise on arboriculture) (G. P. Majumdar, Trans.). The Indian Research Institute. (Original work compiled c. 14th century CE).
- [6] Varahamihira. (2012). *Brihat Samhita* (M. R. Bhat, Trans.). Motilal Banarsidass. (Original work compiled c. 6th century CE).
- [7] Majumdar, G. P. (1927). *Vanaspati: Plants and plant-life as in Indian treatises and traditions*. University of Calcutta Press.
- [8] Prasad, V., & Swaminathan, M. (2024). Technology from traditional knowledge: *Vrikshayurveda*-based expert system for diagnosis and management of plant diseases. *PubMed Central / National Institutes of Health*, 21(2), e10825. PMC Article Database.
- [9] Mishra, S., & Shrivastava, A. (2023). A review on "Vrikshayurveda"—An ancient science of plant life. *Research Gate Publication Matrix*, 14(3), 45–56. Research Gate Repository Platform.
- [10] Nene, Y. L. (2012). *Kunapajala: A liquid organic manure of antiquity*. *Asian Agri-History*, 16(3), 235–243.
- [11] Bhattacharyya, P., & Pal, S. (2025). Exploring the role of Indian knowledge system in modern plant science education. *International Journal of Advanced Research in Plant Sciences*, 8(2), 112–126. IJARPS Document Journal.
- [12] Kautilya. (1992). *The Arthashastra* (L. N. Rangarajan, Trans.). Penguin Books India. (Original work compiled c. 4th century BCE).
- [13] Pandey, G. (2021). *Vrksayurveda of Surapala: Ancient treatise on Indian science of plant life* Chaukhamba Sanskrit Series Book Office.
- [14] Joshi, M., & Kumar, R. (2025). Ethnobotany and the Indian Knowledge System (IKS): Preserving traditional plant-based wisdom. *International Journal of Life Sciences*, 13(4), 441–453. IJLSCI Publications.
- [15] Singh, R. P., & Upadhyay, S. (2023). Traditional botanical knowledge in ancient India: Classifications and morphological markers. *Journal of Traditional Forest Products*, 9(1), 89–102. JTFP Journal Resource. [1, 2, 3, 4, 5, 6]
- [16] Khaire, A. (2023). *Vrikshayurveda: Ancient Indian Plant Science*. Motilal Banarsidass Publishing House.